



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Porcine circovirus type 2 (PCV-2) coinfections in US field cases of postweaning multisystemic wasting syndrome (PMWS)

Citation for published version:

Pallarés, FJ, Halbur, PG, Opriessnig, T, Sorden, SD, Villar, D, Janke, BH, Yaeger, MJ, Larson, DJ, Schwartz, KJ, Yoon, KJ & Hoffman, LJ 2002, 'Porcine circovirus type 2 (PCV-2) coinfections in US field cases of postweaning multisystemic wasting syndrome (PMWS)', *Journal of Veterinary Diagnostic Investigation*, vol. 14, no. 6, pp. 515-519. <https://doi.org/10.1177/104063870201400614>

Digital Object Identifier (DOI):

[10.1177/104063870201400614](https://doi.org/10.1177/104063870201400614)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Veterinary Diagnostic Investigation

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Journal of Veterinary Diagnostic Investigation

<http://vdi.sagepub.com/>

Porcine Circovirus Type 2 (PCV-2) Coinfections in US Field Cases of Postweaning Multisystemic Wasting Syndrome (PMWS)

F. J. Pallarés, P. G. Halbur, T. Opriessnig, S. D. Sorden, D. Villar, B. H. Janke, M. J. Yaeger, D. J. Larson, K. J. Schwartz, K. J. Yoon and L. J. Hoffman

J VET Diagn Invest 2002 14: 515

DOI: 10.1177/104063870201400614

The online version of this article can be found at:

<http://vdi.sagepub.com/content/14/6/515>

Published by:



<http://www.sagepublications.com>

On behalf of:



Official Publication of the American Association of Veterinary Laboratory Diagnosticians, Inc.

Additional services and information for *Journal of Veterinary Diagnostic Investigation* can be found at:

Email Alerts: <http://vdi.sagepub.com/cgi/alerts>

Subscriptions: <http://vdi.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Citations: <http://vdi.sagepub.com/content/14/6/515.refs.html>

>> [Version of Record](#) - Nov 1, 2002

[What is This?](#)

described in other species with lipid mobilization disorders. As with the affected bison in the present study, fatal hepatic lipidosis has been recently described in obese bison in late gestation; therefore, obesity during late gestation should also be included as a key factor in the development of fatal hepatic lipidosis in bison (Berezowski J, Simko E, Feist M, Haigh J, Woodbury M: 2001, Fat bison syndrome: a report of fatal hepatic lipidosis in obese, late pregnant bison (*Bison bison*) cows. *Proc Am Assoc Vet Lab Diagn*: 17.)

A disease syndrome associated with excessive lipid mobilization and hepatic lipidosis has not been previously identified in bison. These findings suggest that pregnant bison are susceptible to such a syndrome. Although the bison in the current report were exposed to various unique stressors that are not present in commercial bison captive herds, bison producers and veterinarians should be aware of the potential for such changes and monitor feeding of pregnant bison as well as potential stressors that could initiate anorexia and negative energy balance situations.

Acknowledgements. The authors thank Dennis Weuve, Larry Wright, Don Robinson, and Terry Krausman for excellent animal care and Aileen Duit, Judi Stasko, Duane Zimmerman, and Derrel Hoy for excellent technical assistance.

Sources and manufacturers

- a. Colorado Serum Co., Denver, CO.

References

1. Bell AW: 1995, Regulation of organic nutrient metabolism during transition from late pregnancy to early lactation. *J Anim Sci* 73: 2804–2819.
2. Cebra CK, Garry FB, Getzy DM, Fettman MJ: 1997, Hepatic lipidosis in anorectic, lactating Holstein cattle: a retrospective study of serum biochemical abnormalities. *J Vet Intern Med* 11: 231–237.
3. Mogg TD, Palmer JE: 1995, Hyperlipidemia, hyperlipemia, and hepatic lipidosis in American miniature horses: 23 cases (1990–1994). *J Am Vet Med Assoc* 207:604–607.
4. Rook JS: 2000, Pregnancy toxemia of ewes, does, and beef cows. *In: Metabolic disorders of ruminants. Veterinary clinics of North America, food animal practice*, vol. 16, ed, TH Herdt, pp. 293–317. WB Saunders, Philadelphia, PA.
5. Rukkwamsuk T, Kruip TAM, Wensing T: 1999, Relationship between overfeeding and overconditioning in the dry period and the problems of high producing dairy cows during the postparturient period. *Vet Q* 21:71–77.
6. Van Saun RJ: 2000, Pregnancy toxemia in a flock of sheep. *J Am Vet Med Assoc* 217:1536–1539.

J Vet Diagn Invest 14:515–519 (2002)

Porcine circovirus type 2 (PCV-2) coinfections in US field cases of postweaning multisystemic wasting syndrome (PMWS)

F. J. Pallarés, P. G. Halbur, T. Opriessnig, S. D. Sorden, D. Villar, B. H. Janke, M. J. Yaeger, D. J. Larson, K. J. Schwartz, K. J. Yoon, L. J. Hoffman

Abstract. The prevalence of different pathogens detected in combination with porcine circovirus type 2 (PCV-2) was studied retrospectively in field cases of postweaning multisystemic wasting syndrome (PMWS) diagnosed at the Iowa State University Veterinary Diagnostic Laboratory, Ames, Iowa, between January 2000, and September 2001. The presence of PCV-2 antigen in lymphoid tissues and/or lung, demonstrated by immunohistochemistry, together with moderate to severe lymphoid depletion and/or granulomatous lymphadenitis, was used as the criteria for the diagnosis of PMWS. A total of 484 cases fulfilled these criteria. Most of the cases (294/369) of PMWS occurred in pigs between the ages of 8 and 18 weeks, with a peak at 10 weeks of age. Porcine reproductive and respiratory syndrome virus was detected in 51.9% of the cases, *Mycoplasma hyopneumoniae* in 35.5%, bacterial septicemia in 14.0%, bacterial pneumonia in 7.6%, swine influenza virus in 5.4%, and PCV-2 alone in 1.9%. In cases with bacterial septicemia the most frequently isolated pathogen was *Streptococcus suis*. In cases with bacterial pneumonia, *Pasteurella multocida* was the most prevalent.

Postweaning multisystemic wasting syndrome (PMWS) associated with porcine circovirus type 2 (PCV-2) has now become a global swine disease problem.^{1,4,10} PMWS was first

described in Canada in 1991 (Clark EG: 1997, *Proc Am Assoc Swine Pract*: 499–501; Harding JCS: 1997, *Proc Am Assoc Swine Pract*: 503) and has since been reported in many countries worldwide.^{8,17,24,30} The disease primarily affects pigs between the ages of 5 and 18 weeks and is clinically characterized by progressive weight loss, tachypnea, dyspnea, anemia, diarrhea, and jaundice.^{13,31} Characteristic microscopic lesions are lymphoid depletion, granulomatous interstitial pneumonia, hepatitis, and nephritis.^{6,12,18,27} Clinical PMWS was reproduced experimentally in some but not all

From the Veterinary Diagnostic Laboratory, College of Veterinary Medicine, Iowa State University, Ames, IA 50011 (Pallarés, Halbur, Opriessnig, Sorden, Villar, Janke, Yaeger, Larson, Schwartz, Yoon, Hoffman), and *Histología y Anatomía Patológica*, Facultad de Veterinaria, Universidad de Murcia, 30071 Murcia, Spain (Pallarés).

Received for publication December 10, 2001.

VIRAL AND BACTERIAL CO-INFECTIONS IN CASES OF PMWS




Figure 1. Number of cases of PMWS for each of the different combinations of pathogens.

of the cesarean-derived, colostrum-deprived pigs,^{7,16} the gnotobiotic pigs,¹¹ the colostrum-deprived conventional pigs,^{2,18} or the conventional pigs⁶ inoculated with PCV-2. All the gnotobiotic pigs infected with PCV-2 and injected with key-hole limpet hemocyanin in incomplete Freund's adjuvant developed lesions characteristic of PMWS, whereas those given only PCV-2 did not develop any lesions.²⁰ Commonly used vaccines such as bacterins for *Mycoplasma hyopneumoniae* and *Actinobacillus pleuropneumoniae* also showed evidence of enhancement of PCV-2 associated disease and lesions.^{5,22} Most recently, characteristic lymphoid and lung lesions of PMWS were reproduced in specific pathogen-free pigs inoculated with an infectious DNA clone of PCV-2; but the pigs did not exhibit wasting.¹²

Several studies have demonstrated enhanced disease and lesions with experimental coinfection of PCV-2 and other infectious agents. Clinical signs of disease were more severe when pigs were coinfecting with PCV-2 and porcine reproductive and respiratory syndrome virus (PRRSV),^{3,16} and PCV-2 replication was potentiated by PRRSV.^{3,28} PMWS also has been reproduced by coinfection of PCV-2 and porcine parvovirus (PPV) in colostrum-deprived pigs² and gnotobiotic pigs.²¹ The objective of this retrospective survey was to examine which coinfecting agents are most commonly diagnosed with PCV-2 in field cases of PMWS submitted to a US diagnostic laboratory that receives swine cases from all over North America.

The presence of PCV-2 antigen in lymphoid tissues and/or lung, demonstrated by immunohistochemistry (IHC),³² together with moderate to severe lymphoid depletion and/or granulomatous lymphadenitis, was used as the criteria for the diagnosis of PMWS. PMWS was not diagnosed if lymphoid tissues were not submitted with the case. The presence of PRRSV and swine influenza virus (SIV) was confirmed by IHC, polymerase chain reaction (PCR), and/or virus isolation (VI). *Mycoplasma hyopneumoniae* was detected by fluorescent antibody (FA) or IHC, but characteristic microscopic lesions of moderate to severe peribronchiolar and perivascular lymphoid hyperplasia were considered diagnostic of *M. hyopneumoniae*-induced pneumonia despite a neg-

ative FA or IHC test for *M. hyopneumoniae*. The isolation of bacterial agents was performed by routine culture procedures. A Kolmogorov-Smirnov test^a was used to determine if the age of the animals affected with PMWS followed a normal distribution.

A total of 4,688 submissions in which PCV-2 IHC had been performed between January 2000, and September 2001, were reviewed. Of these, 1,751 (37.3%) tested positive for PCV-2, and 484 (10.3%) were diagnosed as PMWS. Approximately 80% of the cases were received from Iowa. The rest, in decreasing order, were from North Carolina, Oklahoma, Minnesota, Indiana, Ohio, Colorado, Illinois, Michigan, Kansas, Missouri, Utah, Wisconsin, and Virginia. Of all the animals that tested positive for PCV-2 antigen by IHC, lymphoid lesions characteristics of PMWS were confirmed in only 484 out of 1,751 (27.6%); but lymphoid tissues were not submitted from many of the cases, and thus PMWS could not be ruled out or confirmed in those cases. The number of cases having each of the different combinations of pathogens is illustrated in Fig. 1. Coinfection of PCV-2 + PRRSV was the most common combination with 164 cases. In decreasing order, other coinfections were PCV-2 + *M. hyopneumoniae* (92 cases), PCV-2 + PRRSV + *M. hyopneumoniae* (77 cases), PCV-2 + bacterial septicemia (68 cases), PCV-2 + bacterial pneumonia (37 cases), and PCV-2 + SIV (13 cases). There were only 9 cases in which singular PCV-2 infection was confirmed without any coinfecting pathogens. The total prevalence of coinfection with PCV-2 was 51.9% for PRRSV, 35.5% for *M. hyopneumoniae*, 14.0% for bacterial septicemia, 7.6% for bacterial pneumonia, 5.4% for SIV, and 1.9% for PCV-2 alone.

The types of bacterial pathogens isolated for each of the combinations of PCV-2 + bacterial septicemia and PCV-2 + bacterial pneumonia are illustrated in Figs. 2, 3. The most common bacterial septicemia was *Streptococcus suis* followed by *Salmonella* sp. Bacterial pneumonias were primarily caused by *Pasteurella multocida*.

Ages were provided in 369 of the 484 cases. Among these cases, PMWS primarily occurred between 8 and 18 weeks of age (294/369), with the highest number of cases occurring




Figure 2. Bacterial pathogens isolated in cases of septicemia + PMWS.

at 10 weeks of age. The corresponding histogram followed a normal distribution curve according to the Kolmogorov–Smirnov test ($P > 0.2$) (Fig. 4). Interestingly, there were 8 cases in which PMWS was diagnosed at ≤ 5 weeks of age. Coinfection with PRRSV was found in 6 of these 8 cases.

In some of the PCV-2–positive cases, PMWS confirmation was not possible if lymphoid tissue was not submitted; so the 27.6% is likely a low estimate of the percentage of PCV-2–infected pigs that develop PMWS. In the present survey the prevalence of PMWS was 10.3% (484 cases of PMWS out of 4,868 cases examined), which is very close to the 8.1% (133 cases of PMWS out of 1,634 cases examined) found in a South Korean study during 1999 and 2000,¹⁹ and the 12.9% (7 cases of PMWS out of 59 cases examined) reported in a Canadian study carried out in 1998.²³

The fact that many different infectious agents were isolated in cases of PMWS strongly supports the idea that a variety of pathogens may share a common mechanism in stimulating the immune system,^{5,20} with subsequent progression of PCV-2 infection to PMWS. Alternatively, PCV-2

may make pigs more susceptible to other pathogens. The pathogens commonly involved in porcine respiratory disease complex (PRDC), PRRSV, and *M. hyopneumoniae* were the most common agents that coexisted with PCV-2 in cases of PMWS. PRRSV + PCV-2 coinfection was found in 52% of the cases of PMWS, which is considerably higher than the 29.3% reported in South Korea,¹⁹ more than double the 20% reported in one Canadian field case study,¹ and less than the 67% of the cases with PCV-2 + PRRSV coinfection reported in another Canadian study.²³ European field studies^{25–27} also confirm that PRRSV is involved in many farms with PMWS.

In a recent survey¹⁵ the prevalence of PCV-2 and other pathogens in cases of PRDC received at the Iowa State University Veterinary Diagnostic Laboratory, Ames, Iowa, in the year 2000 was studied, and PCV-2 + PRRSV was demonstrated in 56% of the cases, PCV-2 + *M. hyopneumoniae* in 19%, and PCV-2 + SIV in 12%. PRRSV was the most prevalent coinfecting agent associated with PCV-2 in swine pneumonia cases, with a percentage very similar to that found in the cases of PMWS in this study. The percentage of PRDC cases with PCV-2 + *M. hyopneumoniae* coinfection




Figure 3. Bacterial respiratory pathogens isolated in cases of PMWS.




Figure 4. Age distribution of PMWS cases (369 out of 484 cases provided the age of the pig).

tion was lower and the percentage of PRDC cases with PCV-2 + SIV coinfection was higher than those observed in PMWS cases in this study.

Porcine parvovirus has been used as a model coinfecting agent with PCV-2 to induce PMWS in experimental studies.^{2,18,21} PPV was demonstrated by IHC and PCR in about 18% of the cases of PMWS in Canada⁹ and by in situ hybridization in 25.6% of the cases of PMWS examined in South Korea.¹⁹ Only one of the cases of PMWS in this survey was positive for PPV, but tests for PPV were conducted in only 8 out of 484 cases (5 by FA and 3 by VI). Increased testing of PMWS cases for PPV coinfection is warranted on the basis of the experimental models that clearly demonstrate enhanced clinical signs and lesions typical of PMWS in PCV-2 + PPV-coinfecting pigs.

The most prevalent bacterial coinfection with PCV-2 in this survey was PCV-2 + *S. suis* in 5.5% of the cases. The combination of PCV-2 + *Haemophilus parasuis* was diagnosed in only 1.6% of the cases; but in a recent South Korean study¹⁹ PCV-2 + *H. parasuis* was the most prevalent bacterial coinfection and was found in 32.3% of the PMWS cases.

It was not possible to determine the vaccination status of the PMWS cases in this survey because the case histories provided often did not include vaccination protocols. There is experimental evidence that common vaccines may trigger PCV-2 infection and cause it to progress toward PMWS and that vaccination could potentiate PCV-2 replication.^{5,22} A recent field trial supports the hypothesis that vaccine or drug-induced immuno-modulation can potentiate the effects of PCV-2 infection,²² but the risk of not using vaccines to minimize the effects of coinfections such as *M. hyopneumoniae* may be greater than the risk of vaccination-induced PMWS.

The age of presentation of PMWS followed a normal distribution and coincides with the data from other reports.^{10,14,19} It is uncommon in the US to find PMWS in animals less than 5 weeks of age. In all the 8 cases of PMWS in pigs ≤ 5

weeks of age presented here, other coinfecting agents such as PRRSV were present in addition to PCV-2. Such findings suggest that these pigs have a lack or early loss of maternal immunity against PCV-2 and/or that coinfecting agents such as PRRSV induced increased replication of PCV-2 and earlier development of PMWS.

Perhaps, the strongest argument for the necessity of a coinfection or other cofactors²⁵ for full development of the pathogenic potential of PCV-2 (whether in PMWS or PRDC) is the low percentage of cases in which PCV-2 is found alone (only 1.9% in this study). In a similar South Korean study¹⁹ singular PCV-2 infection was found in 15% of the PMWS cases. With other established pathogens such as PRRSV³³ or SIV (Janke BH, et al.: 2000, Keeping pace with SIV. A. D. Leman Swine Conference, pp. 6–16), the percentage of cases in which they are the only agents found is much higher (39.8% of the PRRSV cases and 36.9% of the SIV cases).

Proven protocols for management and prevention of PMWS are not yet established. There is no PCV-2 vaccine to date. At this point in time, veterinarians often recommend thorough sanitation between groups with the use of disinfectants effective against PCV-2,²⁹ all-in-all-out pig flow, and controlling farm-specific coinfecting agents that may account for progression of PCV-2 infection to PMWS. This survey confirms that in the US, PRRSV and *M. hyopneumoniae* are the most common coinfections in cases of PCV-2-associated PMWS. At least 2 experimental reports^{3,28} support PRRSV-induced increased replication of PCV-2 and increased incidence of PMWS in PCV-2 + PRRSV-coinfecting pigs. Because more than half the cases of PMWS in our study were associated with PRRSV infection, an effective PRRSV control or eradication program may be important in reducing the incidence of PMWS in US herds.

Sources and manufacturers

a. STATISTICA 5.1, 1997, Statsoft, Inc., Tulsa, OK.

References

- Allan GM, Ellis JA: 2000, Porcine circoviruses: a review. *J Vet Diagn Investig* 12:3–14.
- Allan GM, Kennedy S, McNeilly F, et al.: 1999, Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus. *J Comp Pathol* 121:1–11.
- Allan GM, McNeilly F, Ellis J, et al.: 2000, Experimental infection of colostrum deprived piglets with porcine circovirus (PCV2) and porcine reproductive and respiratory syndrome virus (PRRSV) potentiates PCV2 replication. *Arch Virol* 145: 2421–2429.
- Allan GM, McNeilly F, Kennedy S, et al.: 1998, Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. *J Vet Diagn Investig* 10:3–10.
- Allan GM, McNeilly F, Kennedy S, et al.: 2000, Immunostimulation, PCV-2, and PMWS. *Vet Rec* 147:170–171.
- Balasch M, Segales J, Rosell C, et al.: 1999, Experimental inoculation of conventional pigs with tissue homogenates from pigs with post-weaning multisystemic wasting syndrome. *J Comp Pathol* 121:139–148.
- Bolin SR, Stoffregen WC, Nayar GPS, Hamel AL: 2001, Post-weaning multisystemic wasting syndrome induced after experimental inoculation of cesarean-derived, colostrum-deprived piglets with type 2 porcine circovirus. *J Vet Diagn Investig* 13: 185–194.
- Choi C, Chae C, Clark EG: 2000, Porcine postweaning multisystemic wasting syndrome in Korean pig: detection of porcine circovirus 2 infection by immunohistochemistry and polymerase chain reaction. *J Vet Diagn Investig* 12:151–153.
- Ellis JA, Bratanich A, Clark EG, et al.: 2000, Coinfection by porcine circoviruses and porcine parvovirus in pigs with naturally acquired post weaning multisystemic wasting syndrome. *J Vet Diagn Investig* 12:21–27.
- Ellis JA, Hassard L, Clark EG, et al.: 1998, Isolation of circovirus from lesions of pigs with postweaning multisystemic wasting syndrome. *Can Vet J* 39:44–51.
- Ellis J, Krakowka S, Laimore M, et al.: 1999, Reproduction of lesions of postweaning multisystemic wasting syndrome in gnotobiotic piglets. *J Vet Diagn Investig* 11:3–14.
- Fenaux M, Halbur PG, Haqshenas G, et al.: 2002, Cloned genomic DNA of type-2 porcine circovirus is infectious when injected directly into the liver and lymph nodes of pigs: characterization of clinical disease, virus distribution and pathological lesions. *J Virol* 76:541–551.
- Harding JCS, Clark EG: 1997, Recognizing and diagnosing postweaning multisystemic wasting syndrome (PMWS). *Swine Health Prod* 5:201–203.
- Harding JCS, Clark EG, Stokappe JH, et al.: 1998, Postweaning multisystemic wasting syndrome: epidemiology and clinical presentation. *Swine Health Prod* 6:249–254.
- Harms PA, Halbur PG, Sorden SD: 2002, Three cases of porcine respiratory disease complex associated with porcine circovirus type 2 infection. *J Swine Health Prod* 10:27–30.
- Harms PA, Sorden SD, Halbur PG, et al.: 2001, Experimental reproduction of severe disease in CD/CD pigs concurrently infected with type 2 porcine circovirus and porcine reproductive and respiratory syndrome virus. *Vet Pathol* 38:528–539.
- Kennedy S, Allan G, McNeilly F, et al.: 1998, Porcine circovirus infection in Northern Ireland. *Vet Rec* 142:495–496.
- Kennedy S, Moffett D, McNeilly F, et al.: 2000, Reproduction of lesions of postweaning multisystemic wasting syndrome by infection of conventional pigs with porcine circovirus type 2 alone or in combination with porcine parvovirus. *J Comp Pathol* 122:9–24.
- Kim J, Chung H-K, Jung T, et al.: 2002, Postweaning multisystemic wasting syndrome of pigs in Korea: prevalence, microscopic lesions, and coexisting microorganism. *J Vet Med Sci* 64:57–62.
- Krakowka S, Ellis JA, McNeilly F, et al.: 2001, Activation of the immune system is the pivotal event in the production of wasting disease in pigs infected with porcine circovirus-2 (PCV-2). *Vet Pathol* 38:31–42.
- Krakowka S, Ellis JA, Meehan B, et al.: 2000, Viral wasting syndrome of swine: experimental reproduction of postweaning multisystemic wasting syndrome in gnotobiotic swine by coinfection with porcine circovirus 2 and porcine parvovirus. *Vet Pathol* 37:254–263.
- Kyriakis SC, Saoulidis K, Lekkas S, et al.: 2002, The effects of immuno-modulation on the clinical and pathological expression of postweaning multisystemic wasting syndrome. *J Comp Pathol* 126:38–46.
- Larochelle R, Morin M, Antaya M, Magar R: 1999, Identification and incidence of porcine circovirus in routine field cases in Québec as determined by PCR. *Vet Rec* 145:140–142.
- Madec F, Eveno E, Morvan P, et al.: 2000, Post-weaning multisystemic wasting syndrome (PMWS) in pigs in France: clinical observations from follow-up studies on affected farms. *Livest Prod Sci* 63:223–233.
- Quintana J, Segales J, Rosell C, et al.: 2001, Clinical and pathological observations on pigs with postweaning multisystemic wasting syndrome. *Vet Rec* 149:357–361.
- Rodriguez-Arriola GM, Segales J, Rosell C, et al.: 1999, Aujeszky's disease virus infection concurrent with postweaning multisystemic wasting syndrome in pigs. *Vet Rec* 144:152–153.
- Rosell C, Segales J, Plana-Duran J, et al.: 1999, Pathological, immunohistochemical, and in-situ hybridization studies of natural cases of postweaning multisystemic wasting syndrome (PMWS) in pigs. *J Comp Pathol* 120:59–78.
- Rovira A, Balasch M, Segales J, et al.: 2002, Experimental inoculation of conventional pigs with porcine reproductive and respiratory syndrome virus (PRRSV) and porcine circovirus type-2. *J Virol* 76:3232–3239.
- Royer RL, Nawagitgul P, Halbur PG, Paul PS: 2001, Susceptibility of porcine circovirus type 2 to commercial and laboratory disinfectants. *J Swine Health Prod* 9:281–284.
- Segales J, Sitjar M, Domingo M, et al.: 1997, First report of post-weaning multisystemic wasting syndrome in pigs in Spain. *Vet Rec* 141:600–601.
- Sorden SD: 2000, Update on porcine circovirus and postweaning multisystemic wasting syndrome (PMWS). *Swine Health Prod* 8:133–136.
- Sorden SD, Harms PA, Nawagitgul P, et al.: 1999, Development of a polyclonal-antibody-based immunohistochemical method for the detection of type 2 porcine circovirus in formalin-fixed, paraffin-embedded tissue. *J Vet Diagn Investig* 11:528–530.
- Zeman DH: 1996, Concurrent respiratory infections in 221 cases of PRRS virus pneumonia: 1992–1994. *Swine Health Prod* 4: 143–145.